

Hirota equation as an example of integrable symplectic map*

L. Faddeev^{1,†} and A. Yu. Volkov^{2,‡}

¹ *Saint Petersburg Branch of the Steklov Mathematical Institute*

Fontanka 27, Saint Petersburg 191011, Russia

and Research Institute for Theoretical Physics

P. O. Box 9 (Siltavuorenpenger 20C), SF-00014 University of Helsinki, Finland


² *Physique - Mathématique, Université de Bourgogne*

B.P.138, 21004 Dijon Cédex, France


Abstract. The hamiltonian formalism is developed for the Sine-Gordon model on the space-time light-like lattice, first introduced by Hirota. The evolution operator is explicitly constructed in the quantum variant of the model, the integrability of corresponding classical finite-dimensional system is established.

Introduction

We consider a system of difference equations on a plane lattice shown below.



The variables, attached to the vertices of lattice, are denoted by w . The equations connect the values of w 's belonging to a common elementary square



as follows

$$w_N = w_S \frac{f(w_W)}{f(w_E)}. \quad (1)$$

The equation allows to get w_N if three other w 's are given and so defines an evolution in the vertical direction (from south to north).

If the function $f(w)$ is invertible, so that the equation

$$\tilde{w} = f(w) \quad (2)$$

*To appear in Lett. Math. Phys.

†Supported by the Russian Academy of Sciences and the Academy of Finland

‡On leave of absence from Saint Petersburg Branch of the Steklov Mathematical Institute, Fontanka 27, Saint Petersburg 191011, Russia


has a unique solution

$$w = g(\tilde{w}) \quad (3)$$

then the equation (1) can be rewritten in the form


$$\tilde{w}_w = \tilde{w}_E \frac{g(\tilde{w}_S)}{g(\tilde{w}_N)} \quad (4)$$

and defines the evolution from east to west. In what follows we shall consider only move south-north, defining the time axis as vertical. Then the horizontal saw




can be considered as a “line” of initial data.

It is evident that the initial data, given on a portion of that saw, define a solution on the part of a lattice bounded by the punctured lines:




In this sense we can speak of hyperbolicity of our system, the lines being lightlike and defining the light cone.

Alternatively we can use a kind of dual variables ψ to define the same dynamical system. These variables are attached to the vertices of a “dual” lattice



The connection of ψ 's and w 's is given by



$$w = \frac{\psi_W}{\psi_E} \quad (5)$$

and

$$\psi = \prod w \quad (6)$$

where the product is taken over all w 's on the same horizontal line to the east of ψ . The second formula

needs clarification of the boundary conditions; this will be done in the main text for the lattice periodic in the horizontal (spacial) direction.

In terms of ψ 's the equations take the form

$$\psi_N = \psi_S f\left(\frac{\psi_W}{\psi_E}\right) \quad (7)$$

which is also equivalent to

$$\tilde{w} = \frac{\psi_N}{\psi_S} \quad (8)$$

(see the last picture).


In Section 1 we present a symplectic structure on the space of initial data which is conserved by evolution and construct the generator of this evolution. We shall do the latter in a quantum version of our system where this generator is given by an unitary operator. Then in Section 2 for a concrete choice of the function f

$$f(w) = \frac{1 + \kappa^2 w}{\kappa^2 + w} \quad (9)$$

where κ^2 is a real positive parameter we shall show that our symplectic map is completely integrable. Namely, we shall present the set of independent commuting integrals of motion, as many as the number of degrees of freedom. Finally in Section 3 we shall show the equivalence of our system to that introduced by Hirota [1] as the one approximating the famous sine-Gordon equation. Some general comments are collected in Conclusion. Some results of the present paper appeared already in [2].

1 Symplectic map

We shall consider our system on a spacially periodic lattice of even length $2M$. The variables w on a saw of initial data will be numbered as shown on the picture:



We introduce the following Poisson structure

$$\begin{aligned} \{w_{n-1}, w_n\} &= 2\gamma w_{n-1} w_n & n = 2, 3, \dots, 2M \\ \{w_{2M}, w_1\} &= 2\gamma w_{2M} w_1 \\ \{w_m, w_n\} &= 0 & 1 < |m - n| < 2M - 1. \end{aligned} \quad (10)$$

Here γ is a positive parameter playing the role of a coupling constant. The Jacoby identity is evident as the Poisson matrix becomes constant after passing to logarithms of w 's.

In applications one can encounter different variants for the set of values of the variables w . The most evident are the following ones:

- 1) w real and positive;
- 2) w complex and unimodular, $|w| = 1$.

Poisson brackets are compatible with both cases; there are also other possibilities but in this paper we confine ourselves to the second (compact) form of w . It is thus necessary to require that the function f maps the unit circle on itself.

The Poisson structure is degenerate; the functions


$$\begin{aligned} C_1 &= \prod w_{\text{odd}} \\ C_2 &= \prod w_{\text{even}} \end{aligned} \quad (11)$$

are commuting with all w 's. The submanifolds with fixed values of C_1 and C_2 realize the symplectic leaves of our manifold. The number of degrees of freedom is thus $M - 1$.

The map

$$\mathcal{U} : w \longrightarrow \hat{w}$$

where $\hat{w}$'s are the values of the solution of the equations (1) on a saw $\hat{S}$ shifted by time unit from the initial saw S



preserves the Poisson brackets and the values of invariants C_1 and C_2 . Thus $\mathcal{U}$ defines a symplectic map (canonical transformation). It can be described by means of corresponding generating function. We shall not follow this route here and instead shall describe the unitary evolution operator $\mathbf{U}$ corresponding to this map in the quantized variant of our dynamical problem. The reason for this is, that quantum mechanics being more algebraic, the corresponding formulae are just more simple. One can then get the classical variant by the appropriate limit.

After quantization the variables w become unitary operators with the commutation relations

$$\begin{aligned} w_{n-1}w_n &= q^2 w_n w_{n-1} & n = 2, 3, \dots, 2M \\ w_{2M}w_1 &= q^2 w_1 w_{2M} \\ w_m w_n &= w_n w_m & 1 < |m - n| < 2M - 1. \end{aligned} \quad (12)$$

where

$$q = e^{i\hbar\gamma}$$

$\hbar$ being the Planck constant. This commutation relation passes into the Poisson relation (10) with $\hbar \rightarrow 0$ if we identify as usually $\frac{i}{\hbar}[\ , \] = \{ \ , \ \}$.

The evolution operator is given by

$$\mathbf{U} = \prod r(w_{\text{even}}) \prod r(w_{\text{odd}}) \quad (13)$$

where the function $r(w)$ is defined by the function $f(w)$ through the functional equation

$$\frac{r(qw)}{r(q^{-1}w)} = f(w). \quad (14)$$

There is no ordering of factors problem here as only one variable enters the functional equation while in (13) we took into account that all even w 's commute among themselves, the same for odd ones.

Application of $\mathbf{U}$ to w 's is rather trivial due to the locality of their commutation relations. For even w 's we get

$$\hat{w}_{2n} = \mathbf{U}^{-1} w_{2n} \mathbf{U} = r^{-1}(w_{2n-1}) r^{-1}(w_{2n+1}) w_{2n} r(w_{2n+1}) r(w_{2n-1}) \quad (15)$$

which after using the functional equation (14) gives

$$\hat{w}_{2n} = f(qw_{2n+1}) w_{2n} (f(qw_{2n-1}))^{-1}. \quad (16)$$

For odd w 's the calculation consists of several steps but it is as trivial as above. We get


$$\hat{w}_{2n-1} = \mathbf{U}^{-1} w_{2n-1} \mathbf{U} = f(q\hat{w}_{2n}) w_{2n-1} (f(q\hat{w}_{2n-2}))^{-1}. \quad (17)$$

So the iteration of $\mathbf{U}$ leads to the quantum form of the equations of motion

$$w_N = f(qw_W)w_S(f(qw_E))^{-1} \quad (18)$$

which coincides with (1) in the classical limit $q \rightarrow 1$.

Consider now the alternative picture in the variables ψ . The set of variables ψ_n



will be now quasi-periodic

$$\begin{aligned} \psi_{2n+2M} &= C_1 \psi_{2n} \\ \psi_{2n-1+2M} &= C_2 \psi_{2n-1+2M} \end{aligned} \quad (19)$$

and $2M+2$ independent variables $C_1, C_2, \psi_1, \psi_2, \dots, \psi_{2M}$ ($M+1$ degrees of freedom) have the following system of commutation relations:

$$\begin{aligned} \psi_m \psi_n &= q \psi_n \psi_m \quad \text{if } n-m \text{ odd and } 0 < n-m < 2M \\ \psi_{\text{odd}} C_1 &= q^2 C_1 \psi_{\text{odd}} \\ \psi_{\text{even}} C_2 &= q^2 C_2 \psi_{\text{even}} \end{aligned} \quad (20)$$

$$[\psi_{\text{odd}}, \psi_{\text{odd}}] = [\psi_{\text{even}}, \psi_{\text{even}}] = 0.$$

Introduce the evolution by means of the same operator $\mathbf{U}$ with

$$w_n = \frac{\psi_{n+1}}{\psi_{n-1}}. \quad (21)$$

Of course these w 's define system (12) with C_1 and C_2 as monodromies (11).

The derivation of equations of motion for ψ 's is even simpler than that for w 's because each ψ does not commute with only one w

$$\psi_n w_n = q^2 w_n \psi_n. \quad (22)$$

We get

$$\psi_N = \psi_S f(q^{-1} \frac{\psi_W}{\psi_E}) \quad (23)$$

which passes to (7) in the classical limit $q \rightarrow 1$.

2 Integrability

We consider now the particular case with the function $f(w)$ given by

$$f(w) = \frac{1 + \kappa^2 w}{\kappa^2 + w}. \quad (24)$$

Here κ^2 is a fixed positive parameter. We thus have $|f| = 1$ if $|w| = 1$. The inverse function g is given by the same formula after reflection $\kappa^2 \rightarrow -\kappa^2$.

Of course this function has a very special character and it is not clear *a priori* why it is interesting. Its exclusiveness is due to the following property [3]: the solution of the functional equation

$$\frac{r(\lambda, qw)}{r(\lambda, q^{-1}w)} = \frac{1 + \lambda w}{\lambda + w} \quad (25)$$

which we use in the definition of the evolution operator (13)

$$\mathbf{U} = \prod r(\kappa^2, w_{\text{even}}) \prod r(\kappa^2, w_{\text{odd}}) \quad (26)$$

satisfies the Yang-Baxter equation in the following form: let u and v be a Weyl pair

$$uv = q^2vu, \quad (27)$$

then

$$r(\lambda, u)r(\lambda\mu, v)r(\mu, u) = r(\mu, v)r(\lambda\mu, u)r(\lambda, v). \quad (28)$$

This equation allows to interpret r as a fundamental L -operator of quantum chain [4] and use the machinery of QISM (see e.g. [6]). In particular we can introduce a family of matrix L -operators with operator matrix elements


$$L_{n-\frac{1}{2}}(\lambda) = \begin{pmatrix} \psi_{n-1}^{-\frac{1}{2}}\psi_n^{\frac{1}{2}} & \lambda\psi_{n-1}^{-\frac{1}{2}}\psi_n^{-\frac{1}{2}} \\ \lambda\psi_{n-1}^{\frac{1}{2}}\psi_n^{\frac{1}{2}} & \psi_{n-1}^{\frac{1}{2}}\psi_n^{-\frac{1}{2}} \end{pmatrix}. \quad (29)$$

Due to (25) it is connected with r by the commutation relation [3]

$$r(\frac{\lambda}{\mu}, w_n)L_{n+\frac{1}{2}}(\lambda)L_{n-\frac{1}{2}}(\mu) = L_{n+\frac{1}{2}}(\mu)L_{n-\frac{1}{2}}(\lambda)r(\frac{\lambda}{\mu}, w_n) \quad (30)$$

where presumably $w_n = \frac{\psi_{n+1}}{\psi_{n-1}}$.

One can recognize in L a Lax operator for the Hirota equation. Indeed, the equation (23) could be interpreted as a zero curvature condition for the elementary square of our lattice:



$$L_{W \leftarrow N}(\kappa^{-1}\lambda)L_{N \leftarrow E}(\kappa\lambda) = L_{W \leftarrow S}(\kappa\lambda)L_{S \leftarrow E}(\kappa^{-1}\lambda) \quad (31)$$

where

$$L_{B \leftarrow A}(\lambda) = \begin{pmatrix} \psi_A^{-\frac{1}{2}}\psi_B^{\frac{1}{2}} & \lambda\psi_A^{-\frac{1}{2}}\psi_B^{-\frac{1}{2}} \\ \lambda\psi_A^{\frac{1}{2}}\psi_B^{\frac{1}{2}} & \psi_A^{\frac{1}{2}}\psi_B^{-\frac{1}{2}} \end{pmatrix}. \quad (32)$$

It's basically the same zero curvature representation as the one invented by Hirota [1] but this time it is truly quantum.

Multiplying the L -operators along the initial data saw and adding the additional diagonal factor to take care of the quasiperiodicity of ψ 's (see [7]) we get the monodromy matrix

$$T(\lambda) = L_{2M-\frac{1}{2}}(\kappa\lambda)L_{2M-\frac{3}{2}}(\kappa^{-1}\lambda)\dots L_{\frac{3}{2}}(\kappa\lambda)L_{\frac{1}{2}}(\kappa^{-1}\lambda)C_2^{\sigma_3} \quad (33)$$

with commuting traces $t(\lambda) = \text{tr}T(\lambda)$

$$[t(\lambda), t(\mu)] = 0. \quad (34)$$

The last statement is not quite obvious because the L -operator does not seem to fit into the standard ultralocal scheme of QISM. It's somewhat similar to the L -operator of the quantum Volterra model [3]

but involves highly nonlocal variables ψ . The ultralocality is however easy to recover by glueing our L -operators in pairs

$$\begin{aligned}\mathbf{L}_n(\lambda) &= L_{2n-\frac{1}{2}}(\kappa\lambda)L_{2n-\frac{3}{2}}(\kappa^{-1}\lambda) \\ &= \lambda \begin{pmatrix} \lambda^{-1}w_{2n-1}^{\frac{1}{2}} + \lambda w_{2n-1}^{-\frac{1}{2}} & \psi_{2n-1}^{\frac{1}{2}}(\kappa w_{2n-1}^{\frac{1}{2}} + \kappa^{-1}w_{2n-1}^{-\frac{1}{2}})\psi_{2n-1}^{\frac{1}{2}} \\ \psi_{2n-1}^{-\frac{1}{2}}(\kappa^{-1}w_{2n-1}^{\frac{1}{2}} + \kappa w_{2n-1}^{-\frac{1}{2}})\psi_{2n-1}^{-\frac{1}{2}} & \lambda w_{2n-1}^{\frac{1}{2}} + \lambda^{-1}w_{2n-1}^{-\frac{1}{2}} \end{pmatrix}\end{aligned}\quad (35)$$

$$t(\lambda) = \text{tr}(\mathbf{L}_M(\lambda)\mathbf{L}_{M-1}(\lambda)\dots\mathbf{L}_1(\lambda)C_2^{\sigma_3}).$$

Looking at the commutation relations for the operators involved

$$\begin{aligned}\psi_{2n-1}w_{2n-1} &= q^2w_{2n-1}\psi_{2n-1} \\ [\psi_{2m-1}, w_{2n-1}] &= 0 \quad \text{if } m \neq n \\ [\psi_{2m-1}, \psi_{2n-1}] &= [w_{2m-1}, w_{2n-1}] = 0.\end{aligned}\quad (36)$$

we notice that $\mathbf{L}_n$ is not only ultralocal (i.e. matrix elements of the operators $\mathbf{L}_n$ with different numbers n commute) but it also resembles the L -operator of the general lattice XXZ model (compare also with the lattice sine-Gordon L -operator [5]) and indeed one can easily verify that

$$R(\frac{\lambda}{\mu})\mathbf{L}_n(\lambda) \otimes \mathbf{L}_n(\mu) = \mathbf{L}_n(\mu) \otimes \mathbf{L}_n(\lambda)R(\frac{\lambda}{\mu})\quad (37)$$

where $R(\lambda)$ is the usual 4×4 trigonometric R -matrix [6]. This makes the commutativity of $t(\lambda)$ almost evident while the commutation relation (30) (or the zero curvature equation (31)) allows to establish easily that $t(\lambda)$ is conserved by our dynamics

$$\mathbf{U}^{-1}t(\lambda)\mathbf{U} = t(\lambda).\quad (38)$$

Direct inspection shows that $t(\lambda)$ is a polynomial of λ^2 of degree M


$$t(\lambda) = \sum_{k=0}^M I_k \lambda^{2k}\quad (39)$$

and thus $M+1$ coefficients I_k constitute the number of commuting integrals of motion required for the complete integrability. It remains to establish their functional independence; this can be done by usual methods.

In classical limit these integrals remain nontrivial and their existence proves the integrability of our system.

3 Hirota equation

The equations considered above do not coincide exactly with the Hirota equations. The correspondence can be achieved by means of the following change of variables: on each second diagonal line SW-NE of the (dual) lattice



we are to make change $\psi \rightarrow \psi^{-1}$. If the function $f(w)$ is “odd”

$$f(w^{-1}) = f^{-1}(w) \quad (40)$$

(the function $f(w)$ from Section 2 is “odd” while the function $r(\lambda, w)$ is “even”: $r(\lambda, w^{-1}) = r(\lambda, w)$) then the equations of motion in the new variables (call them χ) take the form

$$\chi_N = \chi_S^{-1} f(q^{-1} \chi_W \chi_E). \quad (41)$$

Instead of w ’s we now introduce u ’s

$$u = \chi_W \chi_E. \quad (42)$$

They are connected exactly in the same way as ψ ’s and χ ’s, namely, u coincides with either w or w^{-1} on alternating diagonals of their lattice. The equations of motion in terms of u ’s read

$$u_N = f(qu_W) u_S^{-1} f(q^{-1} u_E). \quad (43)$$

Of course on a periodic lattice all these changes are possible only if the period is even.

The commutation relations for the initial data and the evolution operator now look as follows:

$$\begin{aligned} u_{2n\pm 1} u_{2n} &= q^2 u_{2n} u_{2n\pm 1} & n = 1, 2, \dots, M-1 \\ u_1 u_{2M} &= q^2 u_{2M} u_1 \\ u_{2M-1} u_{2M} &= q^2 u_{2M} u_{2M-1} \\ u_m u_n &= u_n u_m & 1 < |m - n| < 2M-1. \end{aligned} \quad (44)$$

$$\mathbf{U} = \mathbf{U}_0 \prod r(\kappa^2, u_{\text{even}}) \prod r(\kappa^2, u_{\text{odd}}) \quad (45)$$

where $\mathbf{U}_0$ is a reflection of all operators u_n

$$\mathbf{U}_0^{-1} u_n \mathbf{U}_0 = u_n^{-1}. \quad (46)$$

The commutation relations (44) are compatible with this inversion which ensures the existence of $\mathbf{U}_0$. The conservation laws I_n are even function of w ’s and so are commuting with $\mathbf{U}_0$.

It is instructive to remind how equation (41) turns into the sine-Gordon equation in the continuous limit. Introducing variables φ

$$\chi = e^{i\frac{\varphi}{2}} \quad (47)$$

we come to the Hirota form of the equation (41)

$$\sin \frac{1}{4}(\varphi_W + \varphi_E - \varphi_N - \varphi_S) = \kappa^{-2} \sin \frac{1}{4}(\varphi_W + \varphi_E + \varphi_N + \varphi_S). \quad (48)$$

Suppose that when the lattice spacing Δ goes to zero the variables φ turn into the smooth function $\varphi(t, x)$ and thus

$$\varphi_W + \varphi_E - \varphi_N - \varphi_S \longrightarrow \frac{\Delta^2}{2}(\partial_x^2 \varphi - \partial_t^2 \varphi). \quad (49)$$

If now

$$\frac{1}{\Delta^2 \kappa^2} \longrightarrow \frac{m^2}{4} \quad (50)$$

then for $\varphi(t, x)$ we obtain exactly the sine-Gordon equation

$$\partial_t^2 \varphi - \partial_x^2 \varphi + \frac{m^2}{2} \sin 2\varphi = 0. \quad (51)$$

Of course, this limiting procedure is valid only in classical case. The quantum case requires a more delicate balance between the dimensional parameter Δ and dimensionless κ^2 , which accounts for the mass renormalization.

Conclusion

Depending on the interests of a reader, one can look upon this paper from different points of view. For a specialist in dynamical systems the Hirota model gives a family of examples of symplectic maps of a finite-dimensional phase space. In particular, as was explicitly mentioned in [8] for one degree of freedom we get an integrable deformation

$$x_{n+1} - 2x_n + x_{n-1} = -2i \ln \frac{\kappa^2 + e^{-ix_n}}{\kappa^2 + e^{ix_n}}$$

of the so-called “standard map”

$$x_{n+1} - 2x_n + x_{n-1} = -4k \sin x_n.$$

For related results on symplectic maps see also [9, 10].

For a specialist in quantum field theory the quantized Hirota model defines a lattice regularization of a relativistic local field-theoretical model, namely sine-Gordon model. This gives another example of the relation between alternating chain and relativistic massive models observed earlier in [11, 12]. We give a reasonably explicit construction of the evolution operator $\mathbf{U} = e^{-iH\Delta_t}$ where Δ_t is lattice spacing in time direction and H is the hamiltonian. Inclusion of $\mathbf{U}$ into algebra generated by local Lax operator allows to introduce the algebraic Bethe-Ansatz technique for the diagonalization of $\mathbf{U}$. Here we enter the well developed domain (see [6] and the references therein).

A specialist in exactly solvable plane models of statistical mechanics should see here new way of writing and arranging of formulas known in the theory of Z_n model. It is enough to stress that for q being a root of unity $r(\lambda, w)$ is exactly the Fateev-Zamolodchikov R -matrix [14].

Finally, a reader interested in the infinite-dimensional algebraic structures is to see here fragments of a future theory of the algebra $U_q(\hat{sl}(2))$ [15] with which we connect our hopes for the construction of correlation functions in integrable models of quantum field theory. This program is being developed now by the Kyoto group [16].

Acknowledgements. Discussion at different stages of our work with R. Kashaev, V. Tarasov and M. Semenov-Tian-Shansky were most stimulating and we are grateful to all of them.

References

- [1] R. Hirota, J. Phys. Soc. Jap. 43 (1977) 2079.
- [2] A.Yu. Volkov, L. Faddeev, Theor. Math. Phys. 92 (1992) 207.
- [3] A.Yu. Volkov, Phys. Lett. A167 (1992) 345.
- [4] V. Tarasov, L. Takhtajan, L. Faddeev, Theor. Math. Phys. 57 (1983) 163.
- [5] A. Izergin, V. Korepin, Lett. Math. Phys. 5 (1981) 199.
- [6] L. Faddeev, in: Les Houches Lectures 1982 (Elsevier, Amsterdam, 1984) p. 563.
- [7] L. Faddeev, L. Takhtajan, Hamiltonian methods in the theory of solitons (Springer, Berlin, 1987).
- [8] A. Bobenko, N. Kutz, U. Pinkall, SFB preprint No. 42.
- [9] J. Moser, A.P. Veselov, Comm. Math. Phys. 139 (1991) 217.
- [10] F.W. Nijhoff, H.W. Capel, V.G. Papageorgiu, Phys. Rev. A46 (1992) 2155,
F.W. Nijhoff, H.W. Capel, Phys. Lett. A163 (1992) 49.
- [11] L. Faddeev, N. Reshetikhin, Ann. Phys. 167 (1986) 227.

- [12] C. Destri, H.J. de Vega, Nucl. Phys. B290 (1987) 363, J. Phys. A22 (1989) 1329.
- [13] L. Takhtajan, L. Faddeev, Uspekhi Mat. Nauk 34 (1979) 13.
- [14] V. Fateev, A.B. Zamolodchikov, Phys. Lett. A92 (1982) 37.
- [15] I.B. Frenkel, N.Yu. Reshetikhin, Comm. Math. Phys. 146 (1992) 1.
- [16] M. Idzumi, T. Tokihiro, K. Iohara, M. Jimbo, T. Miwa, T. Nakashima, Int. Jour. Mod. Phys. 8A (1993) 1479.